

Product Information

Name of Kit: ImmunoComb[®] Feline Corona Virus (FCoV) [FIP] Antibody Test Kit

Catalog No: 50FFP201/ 50FFP210

No of Tests: 12 (Standard-size Kit)/ 120 (Lab-size Kit)

Intended Use: The ImmunoComb[®] Feline Corona Virus (FCoV) Antibody Test Kit is designed for measuring serum IgG antibody levels to FCoV in cats, to monitor FCoV infections and assist in the diagnosis of Feline Infectious Peritonitis (FIP).

Diagnostic Method: The ImmunoComb[®] test is based on solid phase "dot"-ELISA technology. FCoV antigen is applied to the lowest spot on a comb-shaped plastic card. (The Comb is the solid phase and has 12 teeth-sufficient for 12 test samples.)

The samples to be tested are mixed with diluent in the first row of wells of a multi-chamber developing plate. The Comb is then incubated with the samples in the developing plate. Specific IgG antibodies from the samples, if present, bind to the antigen at the test spots. The middle spot is a positive reference control and the uppermost spot is an internal control, which confirms that the test has been properly developed.

The Comb is then transferred to a well, where unbound antibodies are washed from the antigen spots. In the next step, the Comb is allowed to react with an anti-cat IgG Alkaline Phosphates conjugate, which will bind to antigen-antibody complexes at the test spots. After 2 more washes, the Comb is moved into the last well, where a color result develops via an enzymatic reaction. The intensity of the color result of test spots corresponds directly to the antibody level in the test sample.

| | A* | B* | C* |
|---------------------|-----|-------|------|
| Specificity: | 83% | 99% | 100% |
| Sensitivity: | 95% | 93.4% | 100% |

A* - Data are as published in the article *Evaluation of an in-practice test for feline coronavirus antibodies*, by Dr. Diane D. Addie

B* - Based on unpublished data (2007) obtained in the virology laboratory at the college of Veterinary Medicine in Auburn university

C* - Based on Specificity & Sensitivity data, obtained in Leeds Veterinary laboratories, UK 1997

Pathophysiology: Up to 70% of cats, worldwide, are exposed to Feline Corona Viruses (FCoV). Infection is transmitted by fecal-oral route; the virus can survive in dried secretions for as long as seven weeks. The risk of infection is higher in catteries and multiple-cat households.

FCoV infection is asymptomatic in the majority of cats. In a small percentage of cases, mild signs such as fever, diarrhea and conjunctivitis can occur. This stage may last for an undefined period and then progress to a severe systemic disease, known as Feline Infectious Peritonitis (FIP). Cats of any age or breed may develop FIP. The disease is most often seen in young (less than 2 years old) pedigrees; many of them have a history of recent stress, such as relocation to a new home, surgery (e.g. neutering) or another illness.

Preferred Method of Diagnosis: Serology (measuring serum antibody levels) is the preferred method for monitoring FCoV infections and may help in the diagnosis of FIP when used together with other clinic pathological data.

Main Applications:

- (1) Cats with FIP typically have high levels of anti FCoV antibodies. Therefore, a sero-negative result to FCoV in an ill cat helps rule out the diagnosis of FIP.
- (2) FCoV antibody testing may be used to screen for the presence of FCoV infections in cats before they are introduced into FCoV-free households or catteries.

Interpretation: The level of antibodies (i.e., antibody titer) is determined according to the intensity of the test color result. Thus, no or a light gray color indicates no (negative) or low level of antibodies. Higher levels of antibodies are indicated by darker color results. For the ImmunoComb® Feline Corona Virus Antibody Test Kit a reference spot on each Comb tooth (middle spot) has been calibrated to develop a distinct gray color. This is the same color that is generated by a medium positive result, which is considered to be the 'cut-off' value of a significant antibody titer.

| CombScale Value | Result | Comments |
|------------------------|-----------------|---|
| 0 | Negative | Rules out infection by and shedding of FCoV. |
| 1-2 | Low positive | Possible early or convalescent stage of FCoV infection; may be non-specific result. |
| 3-4 | Medium positive | Indicates previous or current FCoV infection. |
| 5-6 | High positive | Indicates FCoV infection with high risk for shedding virus; consistent with diagnosis of FIP. |

References:

Addie, D. D. (1998). The diagnosis and prevention of FIP and recent research into feline Coronavirus shedding. *ESVIM Proceedings: 8th Annual Congress of the European Society of Veterinary Internal Medicine*.

Addie, D. D. (2000). Guest editorial: Clustering of feline Coronaviruses in multicat households. *The Veterinary Journal*, **159**, 8-9.

Addie, D. D., McLachlan, S. A., Golder, M., Ramsey, I. and Jarrett, O. (2004). Evaluation of an in-practice test for feline coronavirus antibodies. *Journal of Feline Medicine and Surgery*, **6**, 63-67.

Addie, D. D., McLachlan, S. A. & Oswald, J. (2002). Evaluation of the Feline Coronavirus antibody ImmunoComb. *2nd International FCoV/ FIP Symposium, Glasgow, UK*.

Cave, T. A., Golder, M. C., Simpson, J. and Addie, D. D. (2004). Risk factors for feline coronavirus seropositivity in cats relinquished to a UK rescue charity. *Journal of Feline Medicine and Surgery*, **6**, 53-58.

Kiss, I., Kecskemeti, S., Tanyi, J., Klingeborn, B., & Belak, S. (2000). Prevalence and genetic pattern of feline Coronavirus in urban cat populations. *The Veterinary Journal*, **159**, 64-70.

McReynolds, C. & Macy, D. (1997). Feline Infectious Peritonitis. Part II: Treatment and prevention. *The Compendium*, **October**, 1111-1116.

Weiss, R. C. (1991). The diagnosis and clinical management of Infectious Peritonitis. *Veterinary Medicine*, **March**, 308-319.